



Hampshire Biodiversity
Information Centre
HBIC Ref: 7420

Non-statutory designated sites
(Wader and Brent Goose
Strategy Sites only) within 1km
of the Harts Farm Way Site
Boundary (SU706058):

Legend

1km Search Area
Harts Farm Way Site Boundary

Solent Wader Strategy Sites 2017

Most Recent Classification

Uncertain (for positive use)

Uncertain (for negative use)

Uncertain

No recorded use

Important

Solent Brent Goose Strategy Sites 2017

Most Recent Classification

Uncertain (for positive use)

Uncertain (for negative use)

Uncertain

No recorded use

Positive 2002


Important

Created: 13/02/2018
Scale at A3: 1:9,000



0 50 100 200 300 400 500
Meters

© Crown copyright and database rights 2018 Ordnance Survey 100019180. Use of this data is subject to terms and conditions. You are granted a non-exclusive, royalty free, revocable licence solely to view the Licensed Data for non-commercial purposes for the period during which HIC makes it available. You are not permitted to copy, sub-license, distribute, sell or otherwise make available the Licensed Data to third parties in any form. Third party rights to enforce the terms of this licence shall be reserved to Ordnance Survey



Appendix 2

Figure 1

Phase 1 Habitat Survey Plan

Key:



Site boundary



Target Notes (referred to in the report)



Conifer Trees




Dense Scrub



Hardstanding

Figure 1: Phase 1 Habitat Survey Plan (for illustrative purposes only)



Land at Harts Farm Way (Phase 1), Havant - Phase 1 Habitat Survey, 29 January 2018

Target Notes (TNs) corresponding to the Phase 1 Habitat Survey Plan

TN no.	Description
1	Former industrial area dominated by remnant areas of hardstanding and made-up ground, with a mix of short ephemeral and tall ruderal vegetation growing amongst hardcore and rubble. Species included annual meadow-grass (<i>Poa annua</i>), common couch (<i>Elytrigia repens</i>), red fescue (<i>Festuca rubra</i>), nipplewort (<i>Lapsana communis</i>), crane's-bill (<i>Geranium</i> sp.), spear thistle (<i>Cirsium vulgare</i>), willowherb (<i>Epilobium</i> sp.), wavy bitter-cress (<i>Cardamine flexuosa</i>), greater plantain (<i>Plantago major</i>), Butterfly-bush (<i>Buddleja davidii</i>), common mouse-ear (<i>Cerastium fontanum</i>), common nettle (<i>Urtica dioica</i>), white clover (<i>Trifolium repens</i>), hoary ragwort (<i>Senecio erucifolius</i>), bristly oxtongue (<i>Picris echinoides</i>), common mallow (<i>Malva sylvestris</i>), broad-leaved dock (<i>Rumex obtusifolius</i>), black medick (<i>Medicago lupulina</i>), cleavers (<i>Galium aparine</i>) and teasel (<i>Dipsacus fullonum</i>).
2	A row of conifer trees, Leyland Cypress (<i>Cupressus × leylandii</i>), with ground flora including common nettle, campion, willowherb, broad-leaved dock, wild teasel, ivy (<i>Hedera helix</i>), sedges (<i>Carex</i> sp.), cleavers, hoary ragwort, bristly oxtongue, yarrow (<i>Achillea millefolium</i>) and Yorkshire-fog (<i>Holcus lanatus</i>).
3	Dense patches of bramble (<i>Rubus fruticosus</i> agg.).
4	Piles of brash, from previous tree / scrub removal.

Appendix 3 Photographs



Photograph 1 – view to east



Photograph 2 - view to north



Photograph 3 – view to south, including short ephemeral vegetation and conifers



Photograph 4 – view to north-east



Photograph 5 – view to west & hardstanding



Photograph 6 – view to east & conifers